

SECTION A

**A1.**

- (a) Define what is meant by a *topology* on a set  $X$ .
- (b) Define what is meant by saying that a function  $f: X \rightarrow Y$  between topological spaces is *continuous*. Define what is meant by saying that  $f$  is a *homeomorphism*.
- (c) Prove that the closed disc  $\mathbb{D}^2 = \{x \in \mathbb{R}^2 \mid |x| \leq 1\}$  with the usual topology is homeomorphic to the hemisphere  $\{x = (x_1, x_2, x_3) \in S^2 \mid x_3 \geq 0\}$ .  
[Here  $S^2$  denotes the unit sphere  $\{x \in \mathbb{R}^3 \mid |x| = 1\}$  with the usual topology.]

[10 marks]

**Solution**

- (a) Given a set  $X$ , a *topology* on  $X$  is a collection  $\tau$  of subsets of  $X$  with the following properties:
  - (i)  $\emptyset \in \tau, X \in \tau$ ;
  - (ii) the intersection of any two subsets in  $\tau$  is in  $\tau$ :

$$U_1, U_2 \in \tau \Rightarrow U_1 \cap U_2 \in \tau;$$

- (iii) the union of any collection of subsets in  $\tau$  is in  $\tau$ :

$$U_\lambda \in \tau \text{ for all } \lambda \in \Lambda \Rightarrow \bigcup_{\lambda \in \Lambda} U_\lambda \in \tau.$$

[5 marks, bookwork]

- (b)  $f: X \rightarrow Y$  is *continuous* if

$$V \text{ is open in } Y \Rightarrow f^{-1}(V) \text{ is open in } X$$

[1 marks, bookwork]

A *homeomorphism* is a continuous bijection with continuous inverse.

[2 marks, bookwork]

- (c) A homeomorphism  $f: \{\mathbf{x} \in S^2 \mid x_3 \geq 0\} \rightarrow D^2$  is given by  $f(x_1, x_2, x_3) = (x_1, x_2)$  with inverse  $f^{-1}(y_1, y_2) = (y_1, y_2, \sqrt{1 - y_1^2 - y_2^2})$ .

[2 marks, question set]

[Total: 10 marks]

**A2.**

- (a) Define what is meant by saying that a topological space  $X$  is *path-connected*.
- (b) What is meant by saying the path-connectedness is a *topological property*?
- (c) Prove that path-connectedness is a topological property.
- (d) Prove that

$$\{x \in \mathbb{R}^2 \mid |x - (0, 1)| \leq 1 \text{ or } |x + (0, 1)| \leq 1\} \subset \mathbb{R}^2$$

(with the usual topology) is path-connected.

[10 marks]

**Solution**

- (a) A *path* from  $x_0$  to  $x_1$  in  $X$  is a continuous function  $\sigma: [0, 1] \rightarrow X$  with  $\sigma(0) = x_0$  and  $\sigma(1) = x_1$ .  $X$  is said to be *path-connected* if, for each pair of points  $x_0, x_1 \in X$ , there is a path in  $X$  from  $x_0$  to  $x_1$ .

[3 marks, bookwork]

- (b) Saying that path-connectedness is a *topological property* means that, if  $X \cong Y$  are homeomorphic topological spaces, then  $X$  is path connected if and only if  $Y$  is path-connected.

[1 marks, bookwork]

- (c) To prove this, suppose that  $X$  is path-connected. Then, given two points  $y_0, y_1 \in Y$  let  $x_0, x_1 \in X$  be points such that  $f(x_i) = y_i$  (these points exist since  $f$  is a bijection). Since  $X$  is path-connected there is a path  $\sigma: I \rightarrow X$  such that  $\sigma(0) = x_0$  and  $\sigma(1) = x_1$ . Then  $f \circ \sigma: I \rightarrow Y$  is a path in  $Y$  from  $y_0$  to  $y_1$  (since the composition of continuous maps is continuous). Hence,  $Y$  is path-connected. Conversely, if  $Y$  is path-connected then so is  $X$  by the same argument (interchanging the roles of  $X$  and  $Y$ ).

[3 marks, bookwork]

- (d) The two discs  $X_+ = \{x \in X \mid x \geq 0\}$  and  $X_- = \{x \in X \mid x \leq 0\}$  are path-connected.

Now, since  $X = X_+ \cup X_-$  and  $0 \in X_+ \cap X_-$  one can find a path  $\sigma_x^0$  from every point in  $x \in X$  to 0. By composition and inversion one obtains a path between two arbitrary points  $x_0, x_1$ . Such a path can also be stated directly:

$$\sigma(s) = \begin{cases} (1 - 2s)x_0 & s \leq 1/2 \\ (2s - 1)x_1 & s \geq 1/2. \end{cases}$$

[3 marks, new]

**[Total: 10 marks]**

**A3.**

- (a) Define what is meant by saying that a topological space is *Hausdorff*.
- (b) Determine whether the set  $S = \{a, b, c\}$  with topology  $\tau = \{\emptyset, \{a, c\}, \{b\}, \{a, b, c\}\}$  is Hausdorff.
- (c) Suppose that  $X$  and  $Y$  are topological spaces. Define the *product topology* on the Cartesian product  $X \times Y$ . [It is not necessary to prove that this is a topology.]
- (d) Prove that if  $\Delta \subset X \times X$  is closed in the product topology, then  $X$  is Hausdorff.

[10 marks]

**Solution**

- (a) The topological space  $X$  is *Hausdorff* if, for each distinct pair of points  $x, y \in X$ , there exist open sets  $U$  and  $V$  in  $X$  such that  $x \in U$ ,  $y \in V$  and  $U \cap V = \emptyset$ .

[2 marks, bookwork]

- (b) This space is not Hausdorff because every open subset containing  $a$  also contains  $c$  and so open subsets as required cannot be found for  $x = a$  and  $y = c$ .

[3 marks, bookwork]

- (c) The product topology on  $X \times Y$  has a basis

$$\{U \times V \mid U \text{ open in } X, V \text{ open in } Y\},$$

i.e. the open sets consist of all unions of such sets.

[3 marks, bookwork]

- (d) Assume  $\Delta$  is closed. Hence  $X \times X \setminus \Delta$  is open. By definition of the product topology this means it is a union of open rectangles, i.e. sets of the form  $U \times V \subset X \times X \setminus \Delta$  with  $U$  and  $V$  both open in  $X$ . Consider  $x, y \in X$  with  $x \neq y$  then  $(x, y)$  lies outside the diagonal. Hence, it has to be contained in such a set

$$U \times V \subset X \times X \setminus \Delta.$$

On the one hand this implies that  $x \in U$  and  $y \in V$ . On the other hand  $U \cap V = \emptyset$ , since for  $x \in U \cap V$  one would have  $\Delta \ni (x, x) \in U \times V$ . [2 marks, question set]

[Total: 10 marks]

**A4.**

- (a) Suppose that  $X_1$  is a subspace of a topological space  $X$ . Define what is meant by saying that  $X_1$  is a *retract* of  $X$ .
- (b) Use the functorial properties of the fundamental group to prove that, if  $X_1$  is a retract of  $X$ , then, for any  $x_0 \in X_1$ , the homomorphism induced by the inclusion map

$$i_*: \pi_1(X_1, x_0) \rightarrow \pi_1(X, x_0)$$

is injective.

- (c) Hence prove that  $S^1$  is not a retract of the closed disc  $\mathbb{D}^2$ .  
[You may quote any fundamental groups that you need, without proof.]

[10 marks]

**Solution**

- (a)  $X_1 \subset X$  is a retract of  $X$  when there is a continuous map  $r: X \rightarrow X_1$ , such that  $r(x) = x$  for  $x \in X_1$ . [3 marks, bookwork]
- (b) By the functorial properties we have

$$r_* \circ i_* = (r \circ i)_* = (\text{id}_{X_1})_* = \text{id}_{\pi_1(X_1, x_0)}: \pi_1(X_1, x_0) \rightarrow \pi_1(X_1, x_0).$$

Since the composition of  $r_*$  and  $i_*$  is bijective  $r_*$  must be surjective and  $i_*$  must be injective.

[4 marks, bookwork]

- (c) We have  $\pi_1(S^1, x_0) = \mathbb{Z}$  and  $\pi_1(D^2, x_0) = 1$ , the trivial group. But there is not injective map  $\mathbb{Z} \rightarrow \{1\}$ . Hence,  $S^1$  cannot be a retract of  $D^2$ . [3 marks, bookwork]

[Total: 10 marks]

**SECTION B**

**B5.**

- (a) Suppose that  $q: X \rightarrow Y$  is a surjection from a topological space  $X$  to a set  $Y$ . Define the *quotient topology* on  $Y$  determined by  $q$ . State the *universal property* of the quotient topology.
- (b) Suppose that  $f: X \rightarrow Z$  is a continuous surjection from a compact topological space  $X$  to a Hausdorff topological space  $Z$ . Define an equivalence relation  $\sim$  on  $X$  so that  $f$  induces a bijection  $F: X/\sim \rightarrow Z$  from the identification space  $X/\sim$  of this equivalence relation to  $Z$ . Prove that  $F$  is a homeomorphism. [State clearly any general results which you use.]
- (c) Prove that the quotient space  $[0, 1] \times [0, 1]/\sim$  with  $(0, s) \sim (1, s)$  is homeomorphic to the cylinder  $[0, 1] \times S^1 \subset \mathbb{R}^3$ .

[15 marks]

**Solution**

- (a) Given a topological space  $(X, \tau)$  and a surjection  $q: X \rightarrow Y$  the quotient topology on  $Y$  is given by

$$\{V \subset Y \mid q^{-1}(V) \in \tau\}.$$

The *universal property* of the quotient topology is:  $f: Y \rightarrow Z$  to a topological space  $Z$  is continuous if and only if the composition  $f \circ q: X \rightarrow Z$  is continuous.

[4 marks, bookwork]

- (b) Given a continuous surjection  $f: X \rightarrow Z$ , define an equivalence relation on  $X$  by  $x \sim x' \Leftrightarrow f(x) = f(x')$ . Then we may define  $F: X/\sim \rightarrow Z$  by  $F([x]) = f(x)$ . Since  $[x] = [x'] \Leftrightarrow x \sim x' \Leftrightarrow f(x) = f(x')$  (by the definition of the equivalence relation), the function  $F$  is well-defined. Since  $F([x]) = F([x']) \Leftrightarrow f(x) = f(x') \Leftrightarrow x \sim x'$  (by the definition of the equivalence relation) it follows that  $[x] = [x']$  and  $F$  is injective. Since  $f$  is a surjection,  $y = f(x)$  for some  $x \in X$  and so  $y = F([x])$ . Hence  $F$  is a surjection. This shows that  $F: X/\sim \rightarrow Z$  is a bijection. The map  $F: X/\sim \rightarrow Z$  is continuous by the universal property since  $F \circ q = f$  which is given as continuous, where  $q: X \rightarrow X/\sim$  is the quotient map given by  $q(x) = [x]$ .

The space  $X/\sim = q(X)$  is compact since it is the continuous image of a compact set. Hence  $F$  is a homeomorphism since it is a continuous bijection from a compact space to a Hausdorff space.

[7 marks, bookwork]

- (c) To see this, define a surjection  $f: I^2 \rightarrow I \times S^1$  by  $f(x, y) = (x, \exp(2\pi iy))$  where we think of  $S^1$  as  $\{z \in \mathbb{C} \mid |z| = 1\}$  using the standard identification  $\mathbb{C} \cong \mathbb{R}^2$ . This function is continuous by the universal property of the product topology since the component functions are continuous. Now,  $I \times S^1$  is Hausdorff (a subset of Euclidean space) and  $I \times I$  is compact (a closed and bounded subset of Euclidean space). Now the result follows from (b).

[4 marks, bookwork]

[Total: 15 marks]

**B6.**

- (a) Define what is meant by a *compact subset* of a topological space and by a *compact topological space*.
- (b) Prove that, if  $f: X \rightarrow Y$  is a continuous function of topological spaces and  $K \subset X$  is a compact subset, then  $f(K)$  is a compact subset of  $Y$ .
- (c) Given a non-compact Hausdorff space  $(X, \tau)$  consider the set  $X^* = X \sqcup \{\infty\}$  and the topology

$$\tau^* = \tau \cup \{(X \setminus C) \cup \{\infty\} \mid C \subset X \text{ compact}\}.$$

Show that  $(X^*, \tau^*)$  is compact.

[It is not necessary to prove that  $\tau^*$  is a topology.]

[15 marks]

**Solution**

- (a)  $K \subset X$  is compact if each cover of  $K$  by open subsets of  $X$  has a finite subcover.  
If  $X$  itself is a compact subset then  $X$  is a compact space.

[3 marks, bookwork]

- (b) Suppose that  $\mathcal{F}$  is an open cover for  $f(K)$ . Let  $f^{-1}(\mathcal{F}) = \{f^{-1}(V) \mid V \in \mathcal{F}\}$ . Then  $f^{-1}(\mathcal{F})$  is an over cover for  $K$  since, given  $a \in K$ ,  $f(a) \in f(K)$  so that  $f(a) \in V$  for some  $V \in \mathcal{F}$ . Hence  $a \in f^{-1}(V)$  for some  $V \in \mathcal{F}$ .

Now, since  $K$  is compact,  $f^{-1}(\mathcal{F})$  has a finite subcover for  $K$ ,  $\{f^{-1}(V_1), f^{-1}(V_2), \dots, f^{-1}(V_n)\}$ . Thus, given  $b \in f(K)$ ,  $b = f(a)$  for some  $a \in K$ . Then  $a \in f^{-1}(V_i)$  for some  $i$ ,  $1 \leq i \leq n$ , so that  $b = f(a) \in V_i$ . Hence  $\{V_1, V_2, \dots, V_n\}$  is a finite subcover of  $\mathcal{F}$  for  $f(K)$ .

Hence  $f(K)$  is compact.

[6 marks, bookwork]

- (c) Consider an open cover  $\mathcal{F}$  of  $X^*$ . In order to contain  $\infty$  it has to include at least one open subset  $U_\infty$  of the form  $X \setminus C \cup \{\infty\}$  where  $C \subset X$  is compact. Now,  $\mathcal{F}' = \{U \cap X \mid U \in \mathcal{F}\}$  is an open cover of  $X$  (since  $U$  and  $X$  are open in  $X^*$ ) and hence of  $C$ .

By compactness of  $C$  a finite subcover  $\{U_1 \cap X, \dots, U_m \cap X\} \subset \mathcal{F}'$  suffices to cover  $C$ . But then one has the finite subcover  $\{U_\infty, U_1, \dots, U_m\} \subset \mathcal{F}$ .

[6 marks, exercise set]

[Total: 15 marks]

**B7.**

- (a) Prove that, if the product  $\sigma_0 * \tau_0$  of two paths  $\sigma_0$  and  $\tau_0$  in a topological space  $X$  is defined and the paths  $\sigma_1$  and  $\tau_1$  are homotopic to  $\sigma_0$  and  $\tau_0$  respectively, then the product  $\sigma_1 * \tau_1$  is defined and is homotopic to  $\sigma_0 * \tau_0$ .
- (b) Explain how a continuous function  $f: X \rightarrow Y$  induces a homomorphism  $f_*: \pi_1(X, x_0) \rightarrow \pi_1(X, f(x_0))$ . You should indicate why  $f_*$  is well-defined and why it is a homomorphism.
- (c) Prove that, for topological spaces  $X$  and  $Y$  with points  $x_0 \in X, y_0 \in Y$ , there is an isomorphism of groups

$$\pi_1(X \times Y, (x_0, y_0)) \cong \pi_1(X, x_0) \times \pi_1(Y, y_0).$$

[15 marks]

**Solution**

- (a) Given homotopic paths  $H: \sigma_0 \sim \sigma_1$  and  $K: \tau_0 \sim \tau_1$  such that  $\sigma_0 * \tau_0$  is defined. Then  $\sigma_0(1) = \sigma_1(1) = \tau_0(0) = \tau_1(0)$  and so the product  $\sigma_1 * \tau_1$  is defined.

Suppose that  $H: \sigma_0 \sim \sigma_1$  and  $K: \tau_0 \sim \tau_1$ . Then we may define a homotopy  $L: \sigma_0 * \tau_0 \sim \sigma_1 * \tau_1$  by

$$L(s, t) = \begin{cases} H(2s, t) & \text{for } 0 \leq s \leq 1/2 \text{ and } t \in I, \\ K(2s - 1, t) & \text{for } 1/2 \leq s \leq 1 \text{ and } t \in I. \end{cases}$$

This is well defined since, for  $s = 1/2, H(1, t) = x_1 = K(0, t)$ . In addition,  $L$  is continuous by the Gluing Lemma since  $[0, 1/2] \times I$  and  $[1/2, 1] \times I$  are closed subsets of  $I^2$

[5 marks, bookwork]

- (b) The function  $f_*$  is defined by  $f_*([\sigma]) = [f \circ \sigma]$ . It is well-defined since, if  $[\sigma_0] = [\sigma_1]$  then  $\sigma_0 \sim \sigma_1$  and so there exists a homotopy  $H: \sigma_0 \sim \sigma_1$ . Then  $f \circ H: I^2 \rightarrow Y$  gives a homotopy  $f \circ \sigma_0 \sim f \circ \sigma_1$  and so  $[f \circ \sigma_0] = [f \circ \sigma_1]$ .

To see that  $f_*$  is a homomorphism suppose that  $[\sigma], [\tau] \in \pi_1(X, x_0)$ . Then

$$f_*([\sigma][\tau]) = f_*([\sigma * \tau]) = [f \circ (\sigma * \tau)]$$

and

$$f_*([\sigma])f_*([\tau]) = [f \circ \sigma][f \circ \tau] = [(f \circ \sigma) * (f \circ \tau)]$$

and by writing out the formulae we see that  $f \circ (\sigma * \tau) = (f \circ \sigma) * (f \circ \tau): I \rightarrow Y$ . Hence,  $f_*([\sigma][\tau]) = f_*([\sigma])f_*([\tau])$ .

[5 marks, bookwork]

- (c) Let  $p_1: X \times Y \rightarrow X$  and  $p_2: X \times Y \rightarrow Y$  be the projection maps. The function

$$\pi_1(X \times Y, (x_0, y_0)) \rightarrow \pi_1(X, x_0) \times \pi_1(Y, y_0)$$

given by  $\alpha \mapsto ((p_1)_*(\alpha), (p_2)_*(\alpha))$  is an isomorphism. To see this we write down the inverse. Given a loop  $\sigma_1$  in  $X$  based at  $x_0$  and a loop  $\sigma_2$  in  $Y$  based at  $y_0$  then we may define a loop  $\sigma$


in  $X \times Y$  based at  $(x_0, y_0)$  by  $\sigma(s) = (\sigma_1(s), \sigma_2(s))$ . Then  $([\sigma_1], [\sigma_2]) \mapsto [\sigma]$  is well-defined and provides the necessary inverse. Hence the given function is an isomorphism of groups.

[5 marks, question set]

[Total: 15 marks]

**B8.**

- (a) Define what is meant by the *path-components* of a topological space. [You may assume the definition of a path and properties of paths.]
- (b) Prove that a continuous map of topological spaces  $f: X \rightarrow Y$  induces a map  $f_*: \pi_0(X) \rightarrow \pi_0(Y)$  between the sets of path-components, taking care to prove that your function is well-defined. Prove that if  $f$  is a homeomorphism then  $f_*$  is a bijection.
- (c) A pair of distinct points  $\{p, q\}$  in a path-connected topological space  $X$  is called a cut-pair of type  $n$  when the subspace  $X \setminus \{p, q\}$  has  $n$  path-components. Prove that a homeomorphism  $f: X \rightarrow Y$  induces a bijection between the subsets of cut-pairs of type  $n$  for every  $n \in \mathbb{N}$ .
- (d) Hence show, using cut-pairs of type 3 or otherwise, that no two of the following subspaces of  $\mathbb{R}^2$  with the usual topology are homeomorphic.



[15 marks]

**Solution**

- (a) Define an equivalence relation on  $X$  by  $x \sim x'$  if and only if there is a path in  $X$  from  $x$  to  $x'$ . Then the path-components of  $X$  are the equivalence classes.

[2 marks, bookwork]

- (b) Suppose that  $f: X \rightarrow Y$  is a continuous map. Then this induces a function  $f_*: \pi_0(X) \rightarrow \pi_0(Y)$  by  $f_*([x]) = [f(x)]$ . This is well-defined because  $[x] = [x']$  implies that  $x \sim x'$  so that there is a path  $\sigma: [0, 1] \rightarrow X$  in  $X$  from  $x$  to  $x'$ . Then  $f \circ \sigma: [0, 1] \rightarrow Y$  is a path in  $Y$  from  $f(x)$  to  $f(x')$  and so  $[f(x)] = [f(x')]$ .

[3 marks, bookwork]

If  $f$  is a homeomorphism then  $f_*$  is a bijection since the inverse  $g = f^{-1}: Y \rightarrow X$  induces a function  $g_*: \pi_0(Y) \rightarrow \pi_0(X)$  inverse to  $f_*$  since  $g_*(f_*([x])) = [g(f(x))] = [x]$  and  $f_*(g_*([y])) = [y]$ .

[2 marks, bookwork]



- (c) Suppose that  $f: X \rightarrow Y$  is a homeomorphism and  $\{p, q\}$  is a pair of distinct points in  $X$ . Then  $f$  induces a homeomorphism  $X \setminus \{p, q\} \rightarrow Y \setminus \{f(p), f(q)\}$  and this induces a bijection  $f_*: \pi_0(X \setminus \{p, q\}) \rightarrow \pi_0(Y \setminus \{f(p), f(q)\})$ . Hence  $\{p, q\}$  is a cut-pair of type  $n$  in  $X$  if and only if  $\{f(p), f(q)\}$  is a cut-pair of type  $n$  in  $Y$ .

[3 marks, exercise set]

- (d) In space (i) there are two cut-pairs of type 3 (the intersection points of the line segments and the inner or out circle respectively). In space (ii) there is a unique cut-pair of type 3 (the two points at the ends of the diameter). In space (iii) there are infinitely many cut-pairs of type 3 (picking two arbitrary points on the radial line segments).

[5 marks, new]

[Total: 15 marks]

END OF EXAMINATION PAPER